

[twocolumn,showpacs,aps,prevtex4 graphics epsng -0.5in 9.5in document

Ancilla-assisted quantum process tomography J. B. Altepeter, D. Branning, E. Jeffrey, T. C. Wei, and P. G. Kwiat kwiat@uiuc.edu Dept. of Physics, University of Illinois at Urbana-Champaign, Urbana IL 61801-3080

R. T. ThewGroup of Applied Physics, University of Geneva, 1211 Geneva 4, Switzerland J. L. O'Brien M. A. Nielsen A. G. White andrew@physics.uq.edu.au Center for Quantum Computer Technology and School of Physical Sciences, University of Queensland, QLD 4072, Brisbane, Australia

March 7, 2003 42.50.-p, 32.80.-t, 85.60.Gz

abstract Complete and precise characterization of a quantum dynamical process can be achieved via the method of quantum process tomography. Using a source of correlated photons, we have implemented several methods, each investigating a wide range of processes, e.g., unitary, decohering, and polarizing. One of these methods, ancilla-assisted process tomography (AAPT), makes use of an additional “ancilla system,” and we have theoretically determined the conditions when AAPT is possible. Surprisingly, entanglement is not required. We present data obtained using both separable and entangled input states. The use of entanglement yields superior results, however.